



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Hence, if u be capable of expansion in any series of powers of x , and v in a series of integral powers,

$$D^{\mu}uv = D^{\mu}.0 + \left(D^{\mu}u.v + \mu D^{\mu-1}u.v' + \frac{\mu(\mu-1)}{1.2} D^{\mu-2}u.v'' + \dots \right).$$

The restriction of the nature of the expansion of v to *positive integral* powers of x is attended with this advantage that the coefficient of $x^{m+n-\mu}$ in the series for $D^{\mu}uv$ always consists of a finite number of terms, viz. $n+1$.

Note on a Theorem for Expanding Functions of Functions.

BY EMORY MCCLINTOCK, F.I.A., *Milwaukee, Wisconsin.*

It has just come to my knowledge that the "theorem for expanding functions of functions" lately published by me (Vol. II. p. 348) was essentially anticipated, twenty years earlier, by Mr. S. Roberts, F.R.S. Mr. Roberts's brief, yet sufficient statement (Quarterly Journal, Vol. IV. p. 195) is as follows, putting $\phi x = a_0 + a_1x + a_2x^2 + \dots$, and $\Pi = a_1 \frac{d}{da_0} + 2a_2 \frac{d}{da_1} + 3a_3 \frac{d}{da_2} + \dots$:—

"It will be observed that, since $\Pi^n \phi x = D_x^n \phi x$, the notation of (1) applies generally to a function of ϕx and the differentials of ϕx , and we may write

$$F(\phi, \phi', \phi'', \dots) = \epsilon^{x\Pi} F(a_0, a_1, 2a_2, \dots)."$$

MILWAUKEE, Aug. 3, 1880.